

# Effect of Internet Therapeutic Intervention on A1C Levels in Patients With Type 2 Diabetes Treated With Insulin

HUGH D. TILDESLEY, MD; ADEL B. MAZANDERANI, BSC; STUART A. ROSS, MD

**OBJECTIVE**— To assess the effect of an Internet-based glucose monitoring system (IBGMS) on A1C levels in patients with type 2 diabetes treated with insulin.

**RESEARCH DESIGN AND METHODS**— This trial involved 50 patients randomly assigned to receive either conventional treatment alone or with additional follow-up through an IBGMS for 6 months. Patients randomized to the intervention group uploaded blood glucose readings every 2 weeks to a secure Web site for review and receipt of feedback from their endocrinologist. A1C and laboratory test results were collected at 0, 3, and 6 months.

**RESULTS**— The baseline parameters were not significantly different. Over a 6-month followup, A1C dropped from 8.8 to 7.6% ( $P < 0.001$ ) in the intervention group compared with 8.5 to 8.4% ( $P = 0.51$ ) in the control group.

**CONCLUSIONS**— The use of IBGMS significantly improved A1C levels in patients with type 2 diabetes treated with insulin.

*Diabetes Care* 33:1738–1740, 2010

To access full article:

<http://care.diabetesjournals.org/content/33/8/1738.abstract?sid=42835339-9186-4339-8759-7b6b5caf67fe>